



FOR ANIMAL TREATMENT ONLY

PhiCarb 1000

Active constituent: 1000g/kg Nicarbazin

For the prevention and control of coccidiosis caused by *Eimeria spp.* in broiler chickens.

25kg

DIRECTIONS FOR USE

By law the user must take due care, obtaining expert advice when necessary, to avoid unnecessary pain and distress when using the product other than as directed on the label.

RESTRAINTS:

DO NOT USE in birds that are producing or may in the future produce eggs or egg products which may be used or processed for human consumption.

PRECAUTION

Use in birds greater than 28 days of age may increase susceptibility to heat stress.

DOSAGE AND ADMINISTRATION:

Thoroughly mix 125g of PhiCarb 1000 per tonne of feed to produce a concentration of 125g/t nicarbazin in the finished broiler feed. PhiCarb 1000 must be thoroughly mixed in the feed to assure even distribution. It is recommended that the required quantity of PhiCarb 1000 is pre-blended with 4-5 kg of ground cereal before adding to the total mix.

Feed containing PhiCarb 1000 should be the only ration fed to broiler chickens during treatment. Use continuously for 14 days, or as directed by a veterinarian. **Do not use in layer hens.** Consult a veterinarian for advice relating to coccidia control.

WITHHOLDING PERIODS:

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.

MEAT: Animals producing meat or offal for human consumption must not be sold for slaughter during or within 4 days of the last treatment.

EGGS: DO NOT USE in birds that are producing or may in the future produce eggs or egg products which may be used or processed for human consumption.

SAFETY:

May irritate the eyes and nose and throat. Avoid contact with eyes. Do not inhale dust. When opening the container and mixing wear a disposable dust mask covering mouth and nose if dust is present. Wash hands after use.

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126 / New Zealand 0800 764766

DISPOSAL:

Shake container into medicated feed. Do not dispose of undiluted chemicals on-site. Break, crush, or puncture container and deliver to an approved waste management facility. If an approved waste management facility is not available, bury the punctured containers 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

Store below 30°C (room temperature). Protect from light and moisture.

Batch No:

Expiry:

Registered pursuant to the ACVM Act 1997, No A09893.

See www.foodsafety.govt.nz for registration conditions.

Registered to Phibro Animal Health Pty Limited
Level 22, 205 Queen Street
Auckland, New Zealand
1010



A handwritten signature in blue ink, appearing to be "Andrew G. ...".